Scours Treatment

- Standing but depressed
- Good suckle

Give 2 quarts warm electrolytes

Treat with:

- +/- antibiotics
- +/- banamine
- +/- CD antitoxin
 - Keep warm
 - Drench with electrolytes every 8-12 hours until nursing well

- Down/sternal
- Weak suckle
- Cool extremities
- IV catheter
- Give 2 liters IV fluids at 1 drop per second (1 L per hour)
 Treat with :
- +/- antibiotics
- +/- banamine
- +/- CD antitoxin
 - Flush line with heparin
 - Give 1 liter IV fluids at 1 drop per 1.5 seconds
 - Flush line with heparin
- If suckle reflex improves, give oral electrolytes every 8-12 hours until nursing
- +/- Maintenance fluids at 1 drop per 3 seconds until active/recovered

- Down/lateral
- No suckle
- Dry gums
 - IV catheter
 - Give 2 liters at 1 drop per second (1 L per hour)
- Flush line with heparin
- Give 2 liters IV fluids at 1 drop per 1.5 seconds
- Flush line with heparin
 Treat with :
- +/- antibiotics
 - +/- banamine
- +/- CD antitoxin
 - Continue IV fluids if not standing at 1 drop per 3 seconds
 - Once standing with a string suckle, give electrolytes every 8-12 hours until able to nurse

IV fluids

Either Lactated Ringer's solution OR

Sodium bicarb solution

 4 TBSP baking soda in 1 gallon of distilled water If unable to start IV, give 2 L
 lactated ringers over 15 minutes via intraperitoneal route on right side